

Comment on “Vacuum Rabi Splitting in a Semiconductor Circuit QED System”

In this Comment, we challenge the main claims made by Toida *et al.* [1] and demonstrate that their results do not provide direct evidence of vacuum Rabi splitting or vacuum Rabi oscillations. In contrast to statements made by Toida *et al.*, the two sharp parallel structures in Fig. 3(b) of [1] are not indicative of a coherent quantum mechanical interaction. Instead, as shown in previous work [2,3], they are a result of the resonant interaction between the double quantum dot (DQD) and the resonator at detunings $\pm\epsilon$ corresponding to a crossing of the bare DQD transition frequency and the bare resonator frequency. More importantly, a clear anticrossing, allowing for a claim of the observation of strong coherent interaction of the vacuum-Rabi-type, is not observed. Surprisingly, the frequency range of the data displayed in Fig. 4(a) of [1] is narrower than the suggested interaction rate $2g/(2\pi) = 40(60)$ MHz, which does not even in principle allow the resolution of the vacuum Rabi mode splitting in their data. Instead, the data in Fig. 4(b) of [1], reproduced here in Fig. 1(b), show a small frequency shift of less than 2 MHz due to the dispersive interaction between the DQD and the resonator.

The key signature of strong coherent coupling of the vacuum Rabi type is the observation of a resonant mode-splitting with a pair of clearly identifiable distinct modes separated in frequency by $2g/(2\pi)$ [4,5]. The linewidth of these *two distinct modes* on resonance is $\Gamma = \gamma + \kappa/2$, with the resonator energy decay rate κ and the DQD decoherence rate $\gamma = \gamma_1/2 + \gamma_\phi$ determined by its energy decay γ_1 and pure dephasing rates γ_ϕ [6]. From their measurements Toida *et al.* correctly determine $\kappa/(2\pi) = 8$ MHz. However, the authors extract the linewidth of the data shown in Fig. 4(a) of [1] and claim that the maximum




FIG. 1 (color online). (a) Simulation of transmitted power as a function of drive frequency with the DQD on resonance with the resonator for $(\gamma_1, \gamma_\phi, \gamma)/(2\pi) = (8, 8, 12)$ MHz (red open triangles) and $(200, 200, 300)$ MHz (orange solid dots) and $\kappa = 8$ MHz as indicated by the spectrum calculated with the DQD far detuned from the cavity (blue open squares). (b) Comparison of Fig. 4(b) of [1] with our master equation simulation (solid orange lines) using the parameters of [1] but decoherence rates $\gamma_1/(2\pi) = \gamma_\phi/(2\pi) = 200$ MHz. The asymmetry in the data is due to a change in the decoherence rate with ϵ .

observed value represents an accurate measure of Γ on resonance. This is incorrect, as the above expression for Γ requires a resolved spectral measurement of the two vacuum Rabi modes to be applicable [5]. Toida *et al.* mistakenly solve the expression of Γ for the DQD decoherence rate finding a too small estimate of $\gamma/(2\pi) = 12(25)$ MHz [1] resulting in their unjustified claim of having observed the strong coupling limit with $g > \kappa, \gamma$.

To confirm our claims, we have solved the system’s master equation (see Ref. [2]) to determine the expected transmission spectrum in the low photon number limit with the wrongly estimated parameters of [1] finding a clearly resolved vacuum Rabi mode splitting, see red open triangles in Fig. 1(a). However, the authors of [1] do not present this essential data in their work. In addition, numerical calculations of the frequency shifts and linewidths presented in Figs. 4(b),(c) of [1] are in good agreement with the data by Toida *et al.* only when assuming more than 10 times larger values of $\gamma/2\pi = 300$ MHz with $\gamma_1/(2\pi) = \gamma_\phi/(2\pi) = 200$ MHz than claimed in [1], see Fig. 1(b). With these parameters the vacuum Rabi mode splitting is not resolvable, see Fig. 1(a). As a result, using γ extracted from our analysis, the number of Rabi flops $n_{\text{Rabi}} = 0.07 \ll 1$, the critical photon number $n_0 = 112 \gg 1$, and the critical atom number $N_0 = 12 \gg 1$ all lead to the conclusion that the strong coupling regime is *not* reached in [1].

A. Wallraff,¹ A. Stockklauser,¹ T. Ihn,¹ J.R. Petta,² and A. Blais³

¹Department of Physics
ETH Zurich
CH-8093 Zurich, Switzerland

²Department of Physics
Princeton University
Princeton, New Jersey 08544, USA

³Département de Physique
Université de Sherbrooke
Sherbrooke, Québec J1K 2R1, Canada

Received 21 April 2013; published 12 December 2013

DOI: 10.1103/PhysRevLett.111.249701

PACS numbers: 73.21.La, 03.67.Lx, 73.63.Kv

- [1] H. Toida, T. Nakajima, and S. Komiyama, *Phys. Rev. Lett.* **110**, 066802 (2013).
- [2] T. Frey, P.J. Leek, M. Beck, A. Blais, T. Ihn, K. Ensslin, and A. Wallraff, *Phys. Rev. Lett.* **108**, 046807 (2012).
- [3] K.D. Petersson, L.W. McFaul, M.D. Schroer, M. Jung, J.M. Taylor, A.A. Houck, and J.R. Petta, *Nature (London)* **490**, 380 (2012).
- [4] A. Wallraff, D.I. Schuster, A. Blais, L. Frunzio, R.-S. Huang, J. Majer, S. Kumar, S.M. Girvin, and R.J. Schoelkopf, *Nature (London)* **431**, 162 (2004).
- [5] S. Haroche and J.-M. Raimond, *Exploring the Quantum: Atoms, Cavities, and Photons* (Oxford University Press, Oxford, 2006).
- [6] We note that the expression for Γ stated in Ref. [1] is incorrect.