

Evolution of pseudogap phase under pressure and end point of CDW in Nd-LSCO probed by transport measurements

Amirreza Ataei¹, Adrien Gourgout¹, S. Badoux¹, M-E. Boulanger¹, Anaëlle Legros^{1,2}, Clément Collignon^{1,3}, D. Graf⁴, S. Licciardello⁵, S. Wiedmann⁵, J.-S. Zhou⁶, Qianli Ma⁷, Bruce Gaulin^{7,8}, N. Doiron-Leyraud¹, Louis Taillefer^{1,8}

¹Institut Quantique, Département de Physique & RQMP, Université de Sherbrooke, Sherbrooke, QC J1K 2R1, Canada.

²Service de Physique de l'État Condensé (CEA, CNRS), Université Paris-Saclay, CEA Saclay, Gif-sur-Yvette 91191, France

³Laboratoire de Physique et d'Étude des Matériaux, École Supérieure de Physique et de Chimie Industrielles (CNRS), Paris 75005, France

⁴National High Magnetic Field Laboratory, Florida State University, Tallahassee, FL 32306, USA.


⁵High Field Magnet Laboratory (HFML-EMFL) and Institute for Molecules and Materials, Radboud University, 6525 ED Nijmegen, The Netherlands

⁶Materials Science and Engineering Program/Mechanical Engineering, University of Texas - Austin, Austin, TX 78712, USA.


⁷Department of Physics and Astronomy, McMaster University, Hamilton, Ontario L8S 4M1, Canada

⁸Canadian Institute for Advanced Research, Toronto, ON M5G 1Z8, Canada.

Resistivity and Hall effect



Tuning pseudogap critical point with pressure



Thermoelectric power

